

Base-catalyzed synthesis of 2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones and isolation of intermediates using microwave irradiation

A. Davoodnia*, M. Bakavoli, N. Zareei, N. Tavakoli-Hoseini

Department of Chemistry, School of Sciences, Islamic Azad University,
Mashhad Branch, Mashhad 91735-413, Iran

Received September 13, 2008, Revised October 15, 2008

A simple and fast method for the synthesis of some 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones has been developed *via* base-catalyzed cyclocondensation of ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate with isothiocyanates. The uncyclized intermediates, ethyl 4,5-dimethyl-2-[(substituted carbamothioyl)amino]thiophene-3-carboxylates, were isolated when the reactions were carried out under microwave irradiation. These intermediates subsequently underwent cyclization in *t*-butanol in the presence of potassium *t*-butoxide on heating under reflux to give the desired bicyclic products.

Key words: ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate, isothiocyanates, 2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones, microwave irradiation.

INTRODUCTION

Our interest in thieno[2,3-d]pyrimidine synthesis emerges from the numerous reports on their diverse biological activities [1–10]. Various methods have already been proposed for the synthesis of these compounds and the most general ones involve cyclocondensation of suitably functionalized thiophenes with different electrophiles such as chloroformamide [11], α -substituted acetonitriles [12], formic acid [13], phosgene [14], ethyl chloroformate [14] and guanidine [15]. To the best of our knowledge, base-catalyzed cyclocondensation of ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) with isothiocyanates for the synthesis of 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones (**3a–e**) and the utilization of microwave irradiation for isolation of the intermediates (**2a–e**) has not been reported in the literature.

Prompted by these findings and due to our interest in the synthesis of heterocyclic compounds [16–25] and in continuation of our previous works on the synthesis of thieno[2,3-d]pyrimidine derivatives [26–28], we report here a simple and fast method for the synthesis of 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones (**3a–e**) through cyclocondensation of ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) with isothiocyanates under basic conditions.

RESULTS AND DISCUSSION


The starting material (**1**) was prepared according to the literature method [29]. Cyclocondensation of this compound with isothiocyanates in the presence of potassium *t*-butoxide in *t*-butanol under reflux gave products identified as 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones (**3a–e**). Under these conditions, attempts to isolate the reaction intermediates (**2a–e**) failed when we monitored the course of the reactions carefully (Scheme 1).

Due to our interest in the utilization of microwave irradiation for the synthesis of heterocyclic compounds [30–33], we tried to extend this non-conventional synthetic method for the synthesis of compounds **3a–e**. Therefore, ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) was allowed to interact with isothiocyanates under microwave irradiation under solvent-free conditions at 800 W. During monitoring of the reaction mixture by TLC (CHCl₃:MeOH = 95:5), surprisingly, we observed that unexpected products, with R_F-values different from those expected for compounds **3a–e**, were being formed. During work up and identification, it was established that a condensation and not a cyclocondensation reaction had occurred and the intermediates ethyl 4,5-dimethyl-2-[(substituted carbamothioyl)amino]thiophene-3-carboxylates (**2a–e**) were isolated. The reaction did not proceed to form cyclic products even after prolonged irradiation, but when the latter compounds were heated under reflux for 3 hours in the presence of potassium *t*-butoxide

* To whom all correspondence should be sent:
E-mail: adavoodnia@yahoo.com

in *t*-butanol, cyclization reaction occurred and the cyclic products **3a–e** were obtained (Scheme 1).

The structure of the synthesized compounds was deduced from their spectral and microanalytical data. For example, the ^1H NMR spectrum of **2a** did not show the NH_2 signal of the precursor **1** at δ 5.61 ppm, but instead of it showed two broad signals at δ 9.62 and 11.80 ppm belonging to the NH groups indicating the formation of compound **2a**. The IR spectrum showed the absorption bands at 1651, 3215 and 3299 cm^{-1} for carbonyl and two NH groups respectively. The MS of **2a** showed a molecular ion peak at m/z 286 (M^+) corresponding to the molecular formula $\text{C}_{12}\text{H}_{18}\text{N}_2\text{O}_2\text{S}_2$. This compound gave also satisfactory elemental analysis data (See Experimental).



Scheme 1.

In conclusion, we have developed a facile method for the synthesis of 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones (**3a–e**) through cyclocondensation of ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) with isothiocyanates in *t*-butanol containing potassium *t*-butoxide as a base catalyst. Using microwave irradiation, the uncyclized intermediates **2a–e** were isolated.

EXPERIMENTAL

Melting points were measured on a Stuart Model SMP3 melting point apparatus. The IR spectra were obtained on a 4300 Shimadzu spectrophotometer as KBr disks. The ^1H NMR (100 MHz) spectra were recorded on a Bruker AC 100 spectrometer. The mass spectra were determined on a Shimadzu GCMS 17A instrument. Elemental analysis was performed on a Thermo Finnigan Flash EA micro-analyzer. Reactions were performed in a domestic microwave oven Model LG MS-543XD.

Preparation of 3-substituted-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-ones (3a–e). General Procedure. Method A. To a solution of the ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) (5 mmol) and potassium *t*-butoxide (2 mmol) in *t*-butanol (20 ml), the appropriate isothiocyanate (6 mmol) was added. The reaction mixture was heated under reflux for 6 hours. After the completion of the reaction (monitored by TLC, $\text{CHCl}_3:\text{MeOH} = 93:7$), the solvent was evaporated in vacuum, the residue was dissolved in water (15 ml) and subsequently neutralized by 1 N HCl. The crude product was collected and recrystallized from ethanol to give compounds **3a–e** in 75, 77, 68, 86 and 74% yields, respectively.

Method B. A mixture of ethyl 4,5-dimethyl-2-[(substituted carbamothioyl)amino]thiophene-3-carboxylates (**2a–e**) (3 mmol) and potassium *t*-butoxide (1 mmol) in *t*-butanol (15 ml) was heated under reflux for 3 hours. After the completion of the reaction (monitored by TLC, $\text{CHCl}_3:\text{MeOH} = 93:7$), the solvent was evaporated in vacuum, the residue was dissolved in water (15 ml) and subsequently neutralized by 1 N HCl. The crude product was collected and recrystallized from ethanol to give compounds **3a–e** in 79, 78, 73, 91 and 75% yields, respectively.

3-Ethyl-5,6-dimethyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-one (3a). M.p. 257–259°C; ^1H NMR (DMSO-*d*₆, δ ppm): 1.17 (t, 3H, $J = 7$ Hz, CH₃), 2.24 (s, 6H, 2CH₃), 4.36 (q, 2H, $J = 7$ Hz, CH₂), 13.45 (br, 1H, NH); IRS (KBr disc): ν 1684 (C=O), 3137 cm^{-1} (NH); MS, m/z : 240 (M^+); Analytically calculated for $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_2\text{S}_2$: C 49.97; H 5.03; N 11.66; S 26.68. Found: C 50.24; H 5.21; N 11.35; S 26.91.

5,6-Dimethyl-3-phenyl-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-one (3b). M.p. 325–327°C; ^1H NMR (DMSO-*d*₆, δ ppm): 2.24 (s, 6H, 2CH₃), 7.0–7.6 (m, 5H, phenyl), 13.65 (s br, 1H, NH); IRS (KBr disc): ν 1703 (C=O), 3152 cm^{-1} (NH); MS, m/z : 288 (M^+); Analytically calculated for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2\text{S}_2$: C 58.31; H 4.19; N 9.71; S 22.24. Found: C 58.67; H 3.98; N 9.50; S 22.45.

5,6-Dimethyl-3-(3-methylphenyl)-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-one (3c). M.p. 295–297°C; ^1H NMR (DMSO-*d*₆, δ ppm): 2.25 (s, 3H, CH₃), 2.29 (s, 6H, 2CH₃), 6.85–7.45 (m, 4H, arom-H), 13.61 (br, 1H, NH); IRS (KBr disc): ν 1709 (C=O), 3163 cm^{-1} (NH); MS, m/z : 302 (M^+); Analytically calculated for $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_2\text{S}_2$: C 59.57; H 4.67; N 9.26; S 21.21. Found: C 59.28; H 4.89; N 9.51; S 20.97.

5,6-Dimethyl-3-(4-methylphenyl)-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-one (3d). M.p.

270°C (dec); ¹H NMR (DMSO-d₆, δ ppm): 2.24 (s, 3H, CH₃), 2.28 (s, 3H, CH₃), 2.34 (s, 3H, CH₃), 6.90–7.25 (overlapped doublets, 4H, arom-H), 13.60 (br, 1H, NH); IRS (KBr disc): ν 1703 (C=O), 3155 cm⁻¹ (NH); MS, m/z: 302 (M⁺); Analytically calculated for C₁₅H₁₄N₂O₂S₂: C 59.57; H 4.67; N 9.26; S 21.21. Found: C 59.34; H 4.46; N 9.48; S 21.43.

5,6-Dimethyl-3-(4-nitrophenyl)-2-thioxo-2,3-dihydrothieno[2,3-d]pyrimidin-4(1H)-one (3e). M.p. 340–342°C; ¹H NMR (DMSO-d₆, δ ppm): 2.25 (s, 3H, CH₃), 2.32 (s, 3H, CH₃), 7.40–8.45 (overlapped doublets, 4H, arom-H), 13.68 (s br, 1H, NH); IRS (KBr disc): ν 1707 (C=O), 3170 cm⁻¹ (NH); MS, m/z: 333 (M⁺); Analytically calculated for C₁₄H₁₁N₃O₃S₂: C 50.44; H 3.33; N 12.60; S 19.24. Found: C 50.71; H 3.62; N 12.73; S 19.01.

Preparation of ethyl 4,5-dimethyl-2-[(substituted carbamothioyl)amino]thiophene-3-carboxylates (2a–e). *General Procedure*. A mixture of ethyl 2-amino-4,5-dimethylthiophene-3-carboxylate (**1**) (3 mmol) and the appropriate isothiocyanate (4 mmol) was subjected to microwave irradiation at 800 W for 2–3 min (4–5 times). After the completion of the reaction (monitored by TLC, CHCl₃:MeOH = 95:5) the crude product was recrystallized from ethanol to give compounds **2a–e** in high yields.

Ethyl 2-[(ethylcarbamothioyl)amino]-4,5-dimethylthiophene-3-carboxylate (2a). Time 5×2 min; Yield 75%; m.p. 165–167°C; ¹H NMR (DMSO-d₆, δ ppm): 1.44 (t, 3H, J = 6.5 Hz, CH₃), 1.58 (t, 3H, J = 7 Hz, CH₃), 2.32 (s, 6H, 2CH₃), 4.41 (q, 2H, J = 6.5 Hz, CH₂), 4.52 (q, 2H, J = 7 Hz, CH₂), 9.62 (br, 1H, NH), 11.80 (br, 1H, NH); IRS (KBr disc): ν 1651 (C=O), 3215, 3299 cm⁻¹ (two NH); MS, m/z: 286 (M⁺); Analytically calculated for C₁₂H₁₈N₂O₂S₂: C 50.32; H 6.33; N 9.78; S 22.39. Found: C 50.05; H 6.11; N 10.06; S 22.21.

Ethyl 4,5-dimethyl 2-[(phenylcarbamothioyl)amino]thiophene-3-carboxylate (2b). Time 4×3 min; Yield 90%; m.p. 170–172°C; ¹H NMR (DMSO-d₆, δ ppm): 1.22 (t, 3H, J = 7 Hz, CH₃), 2.17 (s, 6H, 2CH₃), 4.20 (q, 2H, J = 7 Hz, CH₂), 7.0–7.6 (m, 5H, phenyl), 10.91 (s, 1H, NH), 11.79 (s, 1H, NH); IRS (KBr disc): ν 1664 (C=O), 3175, 3282 cm⁻¹ (two NH); MS, m/z: 334 (M⁺); Analytically calculated for C₁₆H₁₈N₂O₂S₂: C 57.46; H 5.42; N 8.38; S 19.17. Found: C 57.78; H 5.19; N 8.64; S 18.98.

Ethyl 4,5-dimethyl-2-[(3-methylphenyl)carbamothioyl]amino}thiophene-3-carboxylate (2c). Time 5×3 min; Yield 84%; m.p. 168–169°C; ¹H NMR (DMSO-d₆, δ ppm): 1.48 (t, 3H, J = 7.5 Hz, CH₃), 2.30 (s, 6H, 2CH₃), 2.47 (s, 3H, CH₃), 4.37 (q, 2H, J = 7.5 Hz, CH₂), 7.1–7.6 (m, 4H, arom-H),

11.12 (s, 1H, NH), 12.10 (s, 1H, NH); IRS (KBr disc): ν 1659 (C=O), 3165, 3180 cm⁻¹ (two NH); MS, m/z: 348 (M⁺); Analytically calculated for C₁₇H₂₀N₂O₂S₂: C 58.59; H 5.78; N 8.04; S 18.40. Found: C 58.94; H 6.01; N 7.81; S 18.59.

Ethyl 4,5-dimethyl 2-[(4-methylphenyl)carbamothioyl]amino}thiophene-3-carboxylate (2d). Time 4×3 min; Yield 89%; m.p. 165–167°C; ¹H NMR (DMSO-d₆, δ ppm): 1.25 (t, 3H, J = 7 Hz, CH₃), 2.18 (s, 6H, 2CH₃), 2.29 (s, 3H, CH₃), 4.11 (q, 2H, J = 7 Hz, CH₂), 7.1–7.5 (overlapped doublets, 4H, arom-H), 10.89 (s, 1H, NH), 11.78 (s, 1H, NH); IRS (KBr disc): ν 1658 (C=O), 3178, 3200 cm⁻¹ (two NH); MS, m/z: 348 (M⁺); Analytically calculated for C₁₇H₂₀N₂O₂S₂: C 58.59; H 5.78; N 8.04; S 18.40. Found: C 58.81; H 5.54; N 7.87; S 18.73.

Ethyl 4,5-dimethyl 2-[(4-nitrophenyl)carbamothioyl]amino}thiophene-3-carboxylate (2e). Time 5×3 min; Yield 78%; m.p. 213–215°C; ¹H NMR (DMSO-d₆, δ ppm): 1.26 (t, 3H, J = 7 Hz, CH₃), 2.22 (s, 6H, 2CH₃), 4.30 (q, 2H, J = 7 Hz, CH₂), 7.7–8.4 (overlapped doublets, 4H, arom-H), 11.61 (s, 1H, NH), 12.08 (s, 1H, NH); IRS (KBr disc): ν 1654 (C=O), 3184, 3205 cm⁻¹ (two NH); MS, m/z: 379 (M⁺); Analytically calculated for C₁₆H₁₇N₃O₄S₂: C 50.64; H 4.52; N 11.07; S 16.90. Found: C 50.35; H 4.77; N 10.79; S 16.62.

REFERENCES

1. U. S. Pathak, S. Singh, J. Padh, *Indian J. Chem. Sec. B.*, **30B**, 618 (1991).
2. I. S. Rathod, A. S. Pillai, V. S. Shirsath, *Indian J. Heterocycl. Chem.*, **10**, 93 (2000).
3. *CA* **95**, 115592y (1981).
4. I. A. Kharizomenova, A. N. Grinev, N. V. Samsonova, E. K. Panisheva, N. V. Kaplina, I. S. Nikolaeva, T. V. Pushkina, G. N. Pershin, *Khim.-Farm. Zh.*, **15**, 40 (1981).
5. M. Perrissin, M. Favre, C. Luu-Duc, F. Bakri-Logeais, F. Huguot, G. Narcisse, *Eur. J. Med. Chem.-Chim. Ther.*, **19**, 420 (1984).
6. A. K. El-Ansary, A. H. Omar, *Bull. Fac. Pharm.*, **39**, 17 (2001).
7. U. S. Pathak, N. V. Gandhi, S. Singh, R. P. Warde, K. S. Jain, *Indian J. Chem. Sec. B.*, **31B**, 223 (1992).
8. *CA* **80**, 108567f (1974).
9. *CA* **80**, 70825y (1974).
10. *CA* **115**, 256224y (1991).
11. I. O. Donkor, H. Li, S. F. Queener, *Eur. J. Med. Chem.*, **38**, 605 (2003).
12. C. J. Shishoo, M. B. Devani, U. S. Pathak, S. Ananthan, V. S. Bhadti, G. V. Ullas, K. S. Jain, I. S. Rathod, D. S. Talati, N. H. Doshi, *J. Heterocycl. Chem.*, **21**, 375 (1984).
13. Z. Csuros, R. Soos, J. Palinkas, I. Bitter, *Acta Chim. (Budapest)*, **68**, 397 (1971).
14. F. Sauter, *Monatsh. Chem.*, **101**, 535 (1970).

15. H. Link, *Helv. Chim. Acta*, **73**, 797 (1990).
16. M. Bakavoli, A. Davoodnia, M. Rahimizadeh, M. M. Heravi, M. Ghassemzadeh, *J. Chem. Res. Synop.*, 178 (2002).
17. M. Bakavoli, A. Davoodnia, M. Rahimizadeh, M. M. Heravi, *Phosphorus, Sulfur, Silicon*, **177**, 2303 (2002).
18. M. Roshani, A. Davoodnia, M. Sh. Hedayat, M. Bakavoli, *Phosphorus, Sulfur, Silicon*, **179**, 1153 (2004).
19. A. Davoodnia, M. Bakavoli, A. Vahedinia, M. Rahimizadeh, M. Roshani, *Heterocycles*, **68**, 801 (2006).
20. A. Davoodnia, R. Zhiani, M. Roshani, M. Bakavoli, M. Bashash, *Phosphorus, Sulfur, Silicon*, **182**, 1219 (2007).
21. A. Davoodnia, M. Momen-Heravi, E. Golshani, M. Bakavoli, L. Dehabadi, *J. Chem. Res.*, 257 (2007).
22. A. Davoodnia, M. Bakavoli, M. Bashash, M. Roshani, R. Zhiani, *Turk. J. Chem.*, **31**, 599 (2007).
23. A. Davoodnia, M. Bakavoli, N. Pooryaghoobi, M. Roshani, *Heterocycl. Commun.*, **13**, 323 (2007).
24. A. Davoodnia, M. Bakavoli, Sh. Mohseni, N. Tavakoli-Hoseini, *Monatsh. Chem.*, **139**, 963 (2008).
25. A. Davoodnia, R. Zhiani, N. Tavakoli-Hoseini, *Monatsh. Chem.*, in press.
26. A. Davoodnia, M. Bakavoli, Gh. Barakouhi, N. Tavakoli-Hoseini, *Chin. Chem. Lett.*, **18**, 1483 (2007).
27. A. Davoodnia, H. Behmadi, A. Zare-Bidaki, M. Bakavoli, N. Tavakoli-Hoseini, *Chin. Chem. Lett.*, **18**, 1163 (2007).
28. A. Davoodnia, H. Eshghi, A. Salavaty, N. Tavakoli-Hoseini, *J. Chem. Res.*, 1 (2008).
29. K. Gewald, *Chem. Ber.*, **98**, 3571 (1965).
30. A. Davoodnia, M. Bakavoli, F. Khorramdelan, M. Roshani, *Indian J. Heterocycl. Chem.*, **16**, 147 (2006).
31. A. Davoodnia, M. Rahimizadeh, Sh. Rivadeh, M. Bakavoli, M. Roshani, *Indian J. Heterocycl. Chem.*, **16**, 151 (2006).
32. A. Davoodnia, M. Roshani, E. Saleh Nadim, M. Bakavoli, N. Tavakoli-Hoseini, *Chin. Chem. Lett.*, **18**, 1327 (2007).
33. A. Davoodnia, M. Roshani, S. H. Malaeke, M. Bakavoli, *Chin. Chem. Lett.*, **19**, 525 (2008).

СИНТЕЗА НА 2-ТИОКСО-2,3-ДИГИДРОТИЕНО[2,3-d]ПИРИМИДИН-4(1H)-ОНИ
ЧРЕЗ БАЗИЧНА КАТАЛИЗА И ИЗОЛИРАНЕ НА МЕЖДИННИ СЪЕДИНЕНИЯ
С ИЗПОЛЗВАНЕ НА МИКРОВЪЛНОВО ОБЛЪЧВАНЕ

А. Давудниа*, М. Бакаволи, Н. Зариеи, Н. Таваколи-Хосейни

Департамент по химия, Училище по науки, Ислямски университет Азад, Отдел Маишад,
Маишад 91735-413, Иран

Постъпила на 13 септември 2008 г.; Преработена на 15 октомври 2008 г.

(Резюме)

Предложен е прост и бърз метод за синтезата на 3-заместени-5,6-диметил-2-тиоксо-2,3-дихидроотиено[2,3-d]пиримидин-4(1H)-они чрез циклокондензация на 2-амино-4,5-диметилтиофен-3-карбоксиетилат с изотиоцианати чрез базична катализа. Нециклизираните междинни продукти 4,5-диметил-2-[(заместени карбаматотиоил)амино]тиофен-3-карбоксилати са изолирани когато реакциите се провеждат при микровълново облъчване. Тези междинни продукти впоследствие циклизират в *t*-бутанол в присъствие на калиев *t*-бутоксид при нагряване с обратен хладник до получаване на желаните бициклични продукти.